STING antibody [MSVA-515M] HistoMAX[™]

Cat. No. GTX04484

Host	Mouse	<mark>Package</mark> 500 μl, 100 μl
Clonality	Monoclonal	
lsotype	lgG2c	
Applications	IHC-P	
Reactivity	Human	

PRODUCT This antibody was validated on 76 different Normal Tissues by IHC-P. Summary Go to Normal Tissue Gallery Go to Normal Tissue Gallery Go to Cancer Tissue Gallery

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
IHC-P	1:100-1:200

Note : Manual staining : Heat-induced antigen retrieval for 5 minutes in an autoclave at 121°C in pH 7.8 Tris-EDTA-based Target Retrieval Solution buffer.

Not tested in other applications.

Product Note


Highly recommended for IHC-P in human tissues.

Properties	
Form	Liquid
Buffer	PBS, 0.05% BSA (Please contact us for PBS only format)
Preservative	0.05% sodium azide
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	0.2 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant fragment (around aa190-290) of human TMEM173 protein
Purification	Protein A/G purified
Conjugation	Unconjugated



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 04 / 11 Page 1 of 2




Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

DATA IMAGES



GTX04484 IHC-P Image

IHC-P analysis of human pancreatic ductal adenocarcinoma (PDAC) tissue using GTX04484 STING antibody [MSVA-515M] HistoMAX[™].

Ductal adenocarcinoma with strong STING positivity of all tumor cells.



GTX04484 IHC-P Image

IHC-P analysis of human adrenal gland tissue using GTX04484 STING antibody [MSVA-515M] HistoMAX[™]. STING staining of endothelial cells and of macrophages STING immunohistochemistry.



GTX04484 IHC-P Image

IHC-P analysis of human mucosa from appendix tissue using GTX04484 STING antibody [MSVA-515M] HistoMAX™.

STING staining of inflammatory cells predominates in the interfollicular zone STING immunohistochemistry.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 04 / 11 Page 2 of 2