

ICAM1 / CD54 antibody [N1C2]

Cat. No. GTX100450

Host	Rabbit
Clonality	Polyclonal
Isotype	IgG
Applications	WB, IHC-P
Reactivity	Human, Rat

References (14)

★★★★★ Review (2)

Package

100 µl, 25 µl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000
IHC-P	1:100-1:1000

Not tested in other applications.

Calculated MW 58 kDa. ([Note](#))**Observed MW (kDa)** 58 or 85~110 kDa. The observed M.W. is based on the following publication. PMID: 24474243

Properties

Form	Liquid
Buffer	0.1M Tris, 0.1M Glycine, 10% Glycerol
Preservative	0.01% Thimerosal
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.
Concentration	1 mg/ml (Please refer to the vial label for the specific concentration.)
Immunogen	Recombinant protein encompassing a sequence within the Extracellular domain of human ICAM1. The exact sequence is proprietary.
Purification	Purified by antigen-affinity chromatography.
Conjugation	Unconjugated


Note

For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.

Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.

For full product information, images and publications, please visit our [website](#).

DATA IMAGES



GTX100450 IHC-P Image

ICAM1 / CD54 antibody [N1C2] detects ICAM1 / CD54 protein at cell membrane by immunohistochemical analysis.


Sample: Paraffin-embedded human lung cancer.

Green: ICAM1 / CD54 stained by ICAM1 / CD54 antibody [N1C2] (GTX100450) diluted at 1:250.

Red: alpha Tubulin, a cytoskeleton marker, stained by alpha Tubulin antibody [GT114] (GTX628802) diluted at 1:1000.

Blue: Fluoroshield with DAPI (GTX30920).


Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX100450 IHC-P Image


Immunohistochemical analysis of paraffin-embedded MDAMB468 xenograft, using ICAM1 / CD54(GTX100450) antibody at 1:500 dilution.

Antigen Retrieval: Citrate buffer, pH 6.0, 15 min



GTX100450 WB Image

Untreated (-) and treated (+) A549 whole cell extracts (30 µg) were separated by 5% SDS-PAGE, and the membrane was blotted with ICAM1 / CD54 antibody [N1C2] (GTX100450) diluted at 1:1000. The HRP-conjugated anti-rabbit IgG antibody (GTX213110-01) was used to detect the primary antibody, and the signal was developed with Trident ECL plus-Enhanced.



GTX100450 WB Image

The data was published in the journal PLoS One in 2015. [PMID: 25675397](https://pubmed.ncbi.nlm.nih.gov/25675397/)



For full product information, images and publications, please visit our [website](http://www.genetex.com).