Hepatitis C virus NS3 protein antibody

Cat. No. GTX131276

Host	Rabbit	References
Clonality	Polyclonal	<mark>Package</mark> 100 μl, 25
lsotype	IgG	_
Applications	WB	
Reactivity	Hepatitis C virus	

ferences (3) ckage 0 μl, 25 μl

Applications

Application Note

*Optimal dilutions/concentrations should be determined by the researcher.

Suggested dilution	Recommended dilution
WB	1:500-1:3000


Not tested in other applications.

Properties			
Form	Liquid		
Buffer	PBS, 20% Glycerol		
Preservative	0.025% ProClin 300		
Storage	Store as concentrated solution. Centrifuge briefly prior to opening vial. For short-term storage (1-2 weeks), store at 4°C. For long-term storage, aliquot and store at -20°C or below. Avoid multiple freeze-thaw cycles.		
Concentration	0.58 mg/ml (Please refer to the vial label for the specific concentration.)		
Immunogen	Recombinant protein encompassing a sequence within the C-terminus region of Hepatitis C virus NS3 protein. (HCV 2a (strain JFH-1)). The exact sequence is proprietary.		
Purification	Purified by antigen-affinity chromatography.		
Conjugation	Unconjugated		
Note	For laboratory research use only. Not for any clinical, therapeutic, or diagnostic use in humans or animals. Not for animal or human consumption.		
	Purchasers shall not, and agree not to enable third parties to, analyze, copy, reverse engineer or otherwise attempt to determine the structure or sequence of the product.		




For full product information, images and publications, please visit our <u>website</u>.

DATA IMAGES



GTX131276 WB Image

Serine protease NS3(HCV virus) antibody detects Serine protease NS3 protein by western blot analysis. Non-transfected (-) and Serine protease NS3-transfected (+) Huh7 whole cell extracts (30µg) were separated by 7.5% SDS-PAGE, and the membrane was blotted with Serine protease NS3(HCV virus) antibody (GTX131276) at a dilution of 1:1000



GTX131276 WB Image

Serine protease NS3 (HCV virus) antibody detects Serine protease NS3 (HCV virus) protein by western blot analysis. Un-infected (-) and infected (+, HCV infection for) Huh7 whole cell extracts (20 µg) were separated by 10% SDS-PAGE, and the membrane was blotted with Serine protease NS3 (HCV virus) antibody (GTX131276) diluted by 1:1000.



For full product information, images and publications, please visit our <u>website</u>.

Date 2025 / 04 / 15 Page 2 of 2